Your logo

GP/ OOH Telephone Triage Sepsis Tool

s

To be applied to all children under 5 years with fever (or recent fever) symptoms

NB there is no systems substitute for clinical experience & acumen, but Red Flag Sepsis
will help with early identification of children with systemic response to infection

1. Are any of the following reported? Give safety netting advice: call 999 if child deteriorates rapidly, or call 111/ arrange to see GP if condition fails to improve or gradually worsens. Signpost parent to available resources as appropriate. 2. Is the history suggestive of infection? 4. Is any ONE Amber Flag present? Tick Not responding to social cues, not smiling Parents very worried Wakes only with prolonged stimulation Significantly decreased activity Having to work hard to breathe Poor feeding in infants Reduced urine output 3. Is ONE Red Flag present? Tick Leg pain No response to social cues Cold feet or hands Doesn't wake if roused or won't stay awake If immunity also impaired treat as Red Flag Sepsis Weak, high pitched or continuous cry Grunting or bleating noises with every breath Finding it much harder to breath than normal Very fast breathing/ obvious 'pauses' in breathing Skin that's very pale, mottled, ashen or blue Rash that doesn't fade when pressed firmly

Red Flag Sepsis!

This is time-critical, immediate action is required!

No wet nappies/ not passed urine in last 18 h

Temperature < 36°C (check 3 times in 10 min)

If under 3 months, temperature ≥ 38°C

Immediate actions:

Review immediately and/or dial 999, arrange blue light transfer

Communication:

Fax a brief, clear handover (including observations and antibiotic allergies where present) to receiving Emergency Department

Ensure Paramedics pre-alert as 'Red Flag Sepsis'