




BIRMINGHAM
ST MARY'S
HOSPICE

WHAT IS DELIRIUM?

A guide for
patients and visitors



What is Delirium?

Delirium is a state of mental confusion that can happen if you become medically unwell. It is also known as an 'acute confusional state'.

Medical problems, surgery and medications can all cause delirium. It often starts suddenly and usually lifts when the condition causing it gets better. It can be frightening – not only for the person who is unwell, but also for those around him or her.

What is it like to have Delirium?

You may:

- Be less aware of what is going on around you.
- Be unsure about where you are or what you are doing there.
- Be unable to follow a conversation or to speak clearly.
- Have vivid dreams, which are often frightening and may carry on when you wake up.
- Hear noises or voices when there is nothing or no one to cause them.
- See people or things which aren't there.
- Worry that other people are trying to harm you.
- Be very agitated or restless, unable to sit still and wandering about.
- Be very slow or sleepy.
- Sleep during the day, but wake up at night

- Have moods that change quickly. You can be frightened, anxious, depressed or irritable.
- Be more confused at some times than at others –often in the evening or at night.

How can I help someone with Delirium?

You can help someone with delirium feel calmer and more in control if you:

- Stay calm.
- Talk to them in short, simple sentences. Check that they have understood you. Repeat things if necessary.
- Remind them of what is happening and how they are doing.
- Remind them of the time and date. Make sure they can see a clock or a calendar.
- Listen to them and reassure them.
- Make sure they have their glasses and hearing aid.
- Help them to eat and drink.
- Try to make sure that someone they know well is with them. This is often most important during the evening, when confusion often gets worse.
- If they are in hospital, bring in some familiar objects from home.
- Have a light on at night so that they can see where they are if they wake up.

How common is it?

About 2 in 10 hospital patients have a period of delirium. Delirium is more common in people who are older, have memory problem, have poor hearing or eyesight or have a terminal illness.

Why does it happen?

The most common causes of delirium are:

- a urine or chest infection
- having a high temperature
- side-effects of medicine like pain killers and steroids
- dehydration, low salt levels, low haemoglobin (anaemia)
- liver or kidney problems
- suddenly stopping drugs or alcohol
- terminal illness
- constipation
- being in an unfamiliar place.

There is often more than one cause – and sometimes the cause is not found.

How is delirium treated?

If someone suddenly becomes confused, they need to see a doctor urgently. The person with delirium may be too confused to describe what has happened to them, so it's important that the doctor can talk to someone who knows the patient well.

To treat delirium, you need to treat the cause. For example, an infection may be treated with antibiotics.

Can sedative medication (tranquilisers) help?

Sedatives can make delirium worse, so should only be used in a few situations:

- To calm someone enough to have investigations or treatment.

- To stop someone endangering themselves or other people.
- When someone is very agitated or anxious.
- When someone is seeing or hearing things that are not there.

Low doses of antipsychotic medication should be given to help with frightening hallucinations or beliefs that people are trying to harm you.

Any sedative medication should be given in the lowest possible dose for the shortest time possible.

How long does it take to get better?

Delirium gets better when the cause is treated. You can recover very quickly, but it can take several days or weeks. People with dementia can take a particularly long time to get over delirium.

How do you feel afterwards?

You may not remember what has happened, particularly if you had memory problems beforehand. However, you may be left with unpleasant and frightening memories – and even worry that you are going mad.

It can be helpful to sit down with someone who can explain what happened. This might be a family member, a carer or your doctor. They can go through a diary of what happened each day.

Most people feel relieved when they understand what happened and why.

Will it happen again?

You are more likely to have delirium again if you become medically unwell. Someone needs to keep an eye out for the warning signs that you are getting unwell again – whatever the original cause was. If they are worried they should get a doctor as soon as possible. If medical problems are treated early, this can prevent delirium from happening again.

This information has been taken from the Royal College of Psychiatrists. The factsheet was produced by the RCPsych's Public Education Editorial Board. Series Editor: Dr Philip Timms.

Original authors: Dr Hilary Gordon, Professor George Ikkos, Dr Susie Lingwood and Dr Jim Bolton.

Birmingham St Mary's Hospice

176 Raddlebarn Road Selly Park,
Birmingham, B29 7DA

Tel: 0121 752 8798 / 8799

Fax: 0121 472 4159

Email: info@birminghamhospice.org.uk

www.birminghamhospice.org.uk

St Mary's Hospice Ltd registered in England
No. 1161308. Registered Charity No. 503456